

Holt Physics

Problem 17B**PROBLEM**

Consider three point charges on the x -axis: $q_1 = 4.92 \times 10^{-9} \text{ C}$ is at the origin, $q_2 = -6.99 \times 10^{-8} \text{ C}$ is at $x = -3.60 \times 10^{-1} \text{ m}$, and $q_3 = 5.65 \times 10^{-9} \text{ C}$ is at $x = 1.44 \text{ m}$. Find the magnitude and direction of the resultant force on q_1 .

SOLUTION

Given:

$$q_1 = 4.92 \times 10^{-9} \text{ C} \quad r_{1,2} = -3.60 \times 10^{-1} \text{ m}$$

$$q_2 = -6.99 \times 10^{-8} \text{ C} \quad r_{1,3} = 1.44 \text{ m}$$

$$q_3 = 5.65 \times 10^{-9} \text{ C} \quad k_C = 8.99 \times 10^9 \text{ N}\cdot\text{m}^2/\text{C}^2$$

Unknown: $F_{1,\text{tot}} = ?$

Calculate the magnitude of the forces with Coulomb's law:

$$F_{1,2} = \frac{k_C q_1 q_2}{r_{1,2}^2} = \frac{(8.99 \times 10^9 \text{ N}\cdot\text{m}^2/\text{C}^2)(4.92 \times 10^{-9} \text{ C})(-6.99 \times 10^{-8} \text{ C})}{(-3.60 \times 10^{-1} \text{ m})^2} = -2.39 \times 10^{-5} \text{ N}$$

$$F_{1,3} = \frac{k_C q_1 q_3}{r_{1,3}^2} = \frac{(8.99 \times 10^9 \text{ N}\cdot\text{m}^2/\text{C}^2)(4.92 \times 10^{-9} \text{ C})(5.65 \times 10^{-9} \text{ C})}{(1.44 \text{ m})^2} = 1.21 \times 10^{-5} \text{ N}$$

The forces are all along the x -axis, so add up the x -components:

$$F_{1,\text{tot}} = F_{1,2} + F_{1,3} = -2.39 \times 10^{-5} \text{ N} + 1.21 \times 10^{-5} \text{ N} = \boxed{-1.18 \times 10^{-5} \text{ N}}$$

ADDITIONAL PRACTICE

- Suppose four protons were at the corners of a square. The length of each side of the square is $1.52 \times 10^{-9} \text{ m}$. If q_1 is on the upper right corner, calculate the magnitude and direction of the resultant force on q_1 .
- Consider three point charges, $q_1 = 4.50 \text{ C}$, $q_2 = 4.50 \text{ C}$, and $q_3 = 6.30 \text{ C}$, located at the corners of an isosceles triangle. The charges q_1 and q_2 are 5.00 m apart and form the base. The triangle is 3.50 m high, and q_3 is located at the top. Calculate the magnitude and direction of the resultant force on q_3 .
- Imagine three point charges on the corners of a triangle: $q_1 = -9.00 \text{ nC}$ is at the origin, $q_2 = -8.00 \text{ nC}$ is at $x = 2.00 \text{ m}$, and $q_3 = 7.00 \text{ nC}$ is at $y = 3.00 \text{ m}$. Find the magnitude and direction of the resultant force on q_1 .
- Suppose three point charges are on the y -axis: $q_1 = -2.34 \times 10^{-8} \text{ C}$ is at the origin, $q_2 = 4.65 \times 10^{-9} \text{ C}$ is at $y = 0.500 \text{ m}$, and $q_3 = -2.99 \times 10^{-10} \text{ C}$ is at $y = 1.00 \text{ m}$. What is the magnitude and direction of the resultant force on q_1 ?
- Consider four electrons at the corners of a square. Each side of the square is $3.02 \times 10^{-5} \text{ m}$. Find the magnitude and direction of the resultant force on q_3 if it is at the origin.

6. Imagine three point charges at the corners of an isosceles triangle: $q_1 = 2.22 \times 10^{-10} \text{ C}$, $q_2 = 3.33 \times 10^{-9} \text{ C}$, and $q_3 = 4.44 \times 10^{-8} \text{ C}$. The charges q_1 and q_2 are 1.00 m apart and form the triangle's base. The triangle is 0.250 m tall. If q_3 is at the top, what is the magnitude and direction of the resultant force on q_3 ?
7. Consider three 2.0 nC point charges at the following locations: at (0 m, 0 m), at (1.0 m, 2.0 m), and at (1.0 m, 0 m). Find the magnitude and direction of the resultant force on the charge at the origin.
8. Consider three point charges on the corners of a triangle, where $q_1 = -4.0 \text{ mC}$ at the origin; $q_2 = -8.0 \text{ mC}$ at (2.0 m, 0 m); and $q_3 = 2.0 \text{ mC}$ at (0 m, 2.0 m). Calculate the magnitude and direction of the resultant force on q_1 .
9. Suppose three point charges are on the corners of a triangle: $q_1 = 9.00 \text{ mC}$ is at the origin, $q_2 = 6.00 \text{ mC}$ is at the point (1.00 m, 1.00 m), and $q_3 = 3.00 \text{ mC}$ is at (-1.00 m, 1.00 m). Find the magnitude and direction of the resultant force on q_1 .
10. Consider three equal point charges of 4.00 nC on a line. All charges are 4.00 m apart. Calculate the magnitude and direction of the resultant force on the charge in the middle.